

INSECTICIDE RECOMMENDATIONS FOR SMALL GRAINS - 2009
(Barley, Oats, Wheat)

ENT-47

Prepared by D. W. Johnson, Extension Entomologist

Small grains are multi-purpose crops that occupy an important place in production systems. Small grains are attacked by a wide variety of insect pests, but good cultural practices can greatly reduce the potential for economic losses:

- Plant at the proper time (see AGR-18). Planting too early can result in problems with fall armyworms, Hessian flies, aphids, and diseases carried by insects.
- Use recommended seeding rates. Stands that are too thick provide good over-wintering sites for pests and are prone to armyworm infestations in the spring.
- Do not apply excess nitrogen. Luxuriant growth promotes aphid and armyworm problems. To establish need and estimate timing of insect control in small grains, fields should be checked weekly when temperatures are above 50°F from emergence to maturity.
- Know the difference between pest insects and beneficial insects.
- Check with your County Extension Agent for Agriculture about training in pest identification, damage thresholds, and control measures (Integrated Pest Management and Pesticide Safety Education training).

Additional Information

In addition to these recommendations the producer is advised to review **IPM-4 Kentucky IPM Manual for Small Grains**. This publication will provide information about identification, life cycle, scouting techniques, and threshold values for the common pests of small grains. This manual may be found on the IPM web pages at: <http://www.uky.edu/Agriculture/IPM/ipm.htm>.

Additionally, you may find useful information about a specific pest in our ENTFACT series. These fact sheets may be found on the Entomology web pages at: <http://www.uky.edu/Agriculture/Entomology/entfacts.htm>. These and other publications and educational materials are also available to the producer through your County Extension Office.

Use Insecticides Properly

Products listed in this publication are not the only products labeled for use. These products are commonly used and generally available in Kentucky. You may find many other products with different trade names containing the same active ingredient. Be sure the product you choose is labeled for the intended use and registered in Kentucky.

This publication is an abbreviated guide; it is **not a substitute for a product label**. Before using an insecticide, read the entire label. Note sections containing directions for use, and the warning and precautionary statements. Be thoroughly familiar with the proper safety equipment (i.e., goggles, protective suits, respirators, etc.) required to afford maximum protection. Those involved in control operations should always know the name of the chemical being used and the particular concentration being applied.

Chemicals listed in *bold italics* are **Restricted Use** pesticides. Persons buying, using, or supervising the use of these pesticides must be certified as competent to do so. Certification training is available from your county extension agent for agriculture. Check <http://www.uky.edu/Agriculture/PAT/welcome.htm> for information on certification.

Seed Treatment for Aphid Control

Use only on very early planting in fields with a history of barley yellow dwarf virus problems.

The seed applied insecticides Cruiser (thiamethoxam) and Gaucho (imidacloprid) are registered for use on Wheat and Barley. These products must be applied by a commercial seed treater. The manufacturers, Syngenta and Bayer respectively, control the rates and combination of products. See your seed supplier for information.

Post Emergence Treatments for Aphids

(See: Aphids and Barley Yellow Dwarf in Kentucky Grown Wheat. Entfact-121.
<http://www.uky.edu/Agriculture/Entomology/entfacts/pdfs.entfa121.pdf>)

The most important time for controlling aphids to prevent BYD is the first 30 days following emergence. The second most important time is the second 30 days following emergence. Generally, an insecticide applied after the wheat reaches Feeke's 4.0 does little good. Current research data support the following treatment guidelines.

Numbers of aphids per foot of wheat row required to support an insecticide application for management of BYD.

<u>Crop Age</u>	<u>Aphids/Foot of Row</u>
30 days post emergence	3
30 to 60 days post emergence	6
More than 60 days post emergence	10

Delayed planting until after the Hessian Fly free date (Oct. 10 north, to Oct. 15 south) will reduce aphid pressure in the fall. In late spring, treat only if large numbers of aphids are present and plants are exhibiting stress symptoms, or if there are an average of 50 or more aphids per head during the grain filling stage.

Foliar Treatments for Aphids

Insecticide	Rate per Acre	Days to Harvest
<i>Baythroid 2</i> (Wheat only) (cyfluthrin)	1.8 to 2.4 fl. oz.	30 (Grain) 3 (grazing or forage)
<i>Lannate SP</i> (methomyl)	1/4 to 1/2 lb.	7 (Grain) 10 (grazing or feeding)
<i>Mustang Max</i> (Wheat only) (zeta-cypermethrin)	3.2 to 4.0 fl. oz.	14 (grain, forage & hay)
<i>Pennacap-M</i> (methyl parathion)	2 to 3 pts.	14 (Harvest or grazing)
<i>Warrior</i> (Wheat Only) (lambda-cyhalothrin)	2.56 to 3.84 fl. oz.	30 (Grain & Hay)

Foliar Treatments for Armyworms

(See: Armyworms in Small Grains. Entfact-111.
<http://www.uky.edu/Agriculture/Entomology/entfacts/pdfs.entfa111.pdf>)

Insecticide	Rate per Acre	Days to Harvest
Bacillus thuringiensis "B.t."	Aid in control of armyworms can be obtained using many products containing "B.t.". Some examples are Dipel, Javelin and Lepinox. Check label for details. Days to Harvest = 0.	
<i>Baythroid 2</i> (Wheat only) (cyfluthrin)	1.8 to 2.4 fl. oz.	30 (Grain) 3 (grazing or forage)
<i>Lannate SP</i> (methomyl)	1/4 to 2 lbs.	7 (Grain) 10 (Grazing & feeding)
Malathion 5 (malathion)	2 pts.	7
<i>Mustang Max</i> (Wheat Only) (zeta-cypermethrin)	1.76 to 4.0 fl. oz	14 (Grain, forage & hay)
<i>Pennacap M</i> (methyl parathion)	2 to 3 pts.	14 (Harvest or grazing)

Sevin 80 (Wheat Only) (carbaryl)	1-1/4 to 1-7/8 lbs.	21 (Grain or Straw) 7 (Grazing or forage)
Tracer (spinosad)	1.5 to 3.0 fl. oz.	21(Grain or straw) 14(forage or hay)
Warrior (Wheat Only) (lamda-cyhalothrin)	2.56 to 3.84 fl. oz.	30 (Grain & Hay)

Foliar Treatment for Cereal Leaf Beetle

(See: Cereal Leaf Beetle in Kentucky Wheat. Entfact-107.

<http://www.uky.edu/Agriculture/Entomology/entfacts/pdfs.entfa107.pdf>)

Treatment is justified if population size reaches one cereal leaf beetle (adult or larvae) per stem.

Insecticide	Rate per Acre	Days to Harvest
Baythroid 2 (Wheat only) (cyfluthrin)	1.0 to 1.8 fl. oz.	30 (Grain) 3 (grazing or forage)
Lannate SP (methomyl)	¼ to 1/2 lb	7 (Grain) 10 (Grazing & Feeding)
Malathion 5 (malathion)	1 to 1 1/2 pts.	7
MustangMax (Wheat Only) (zeta-cypermethrin)	1.76 to 4.0 fl. oz	14 (Grain, forage & hay)
Sevin 80 WSP (Wheat Only) (carbaryl)	1-1/4 lbs.	21 (Grain or Straw) 7 (Grazing or forage)
Tracer (naturalyte)	1.0 to 3.0 fl. oz.	21(Grain or straw) 14(forage or hay)
Warrior (Wheat Only) (lambda-cyhalothrin)	2.56 to 3.84 fl. oz.	30 (Grain & Hay)

Foliar Treatments for Grasshoppers

(See: Three Common Kentucky Grasshoppers and their Natural Enemies. Entfact-116.

<http://www.uky.edu/Agriculture/Entomology/entfacts/pdfs.entfa116.pdf>)


Insecticide	Rate per Acre	Days to Harvest/Forage
Baythroid 2 (Wheat only) (cyfluthrin)	1.8 to 2.4 fl. oz.	30 (Grain) 3 (grazing or forage)
Dimethoate 4 (Wheat Only) (dimethoate)	3/4 pt.	35 (Grain) 14 (Grazing)
Furadan 4F (carbofuran)	1/4 to 2 pt.	Spray before heads emerge Do not feed to livestock
Mustang Max (Wheat Only) (zeta-cypermethrin)	3.2 to 4.0 fl. oz	14 (Grain, forage & hay)
PennCap M (methyl parathion)	2-3 pts.	14 (Harvest or grazing)
Sevin 80 WSP (Wheat Only) (carbaryl)	2/3 to 1-7/8 lbs.	21 (Grain or Straw) 7 (Grazing or forage)
Warrior (Wheat Only) (lambda-cyhalothrin)	2.56 to 3.84 fl. oz.	30 (Grain & Hay)

HESSIAN FLY

(See: Hessian Fly in Kentucky Wheat. Entfact-101.
<http://www.uky.edu/Agriculture/Entomology/entfacts/pdfs.entfa101.pdf>)

Delay in planting until October 10 (north) or October 15 (south) will control this fly. No chemical treatments are recommended.

Map showing normal safe dates for sowing wheat to escape injury by the Hessian fly. These are approximate and will vary some from year to year.



Products for Control of Insect Pests in Stored Small Grains

(See: Controlling Insect in Stored Grain. Entfact-145.
<http://www.uky.edu/Agriculture/Entomology/entfacts/pdfs.entfa145.pdf>)

The availability of stored grain insecticides is undergoing continuous change. Always check the label of the product to insure that you use it correctly

“Clean-out” Fumigant

Applied to boots of elevators, beneath false floors, etc. This is an “empty” space fumigation targeted at the space beneath the perforated floor in a metal grain bin. Fumigant is applied on a volume not bushel basis. See the **WARNING** below.

Chlor-O-Pic (chloropicrin) 2-4 lbs. / 1000 square feet **Note: Do NOT use chloropicrin to treat grain!**

Chloropicrin is significantly heavier than air and is therefore preferred for “clean-out” fumigations. However, though it is still labeled for “clean-out” of empty bins, it may not be available due to shipping constraints. Chloropicrin is highly corrosive to most metals.

Phostoxin, Fumtoxin etc. (aluminum phosphide) tablets 30-140 / 1000 cubic feet.
 pellets 150-700 / 1000 cubic feet

Note: Applied by volume NOT by bushels.

Aluminum phosphide is not significantly heavier than air and is therefore not the preferred product. However, it is labeled for this use and is easily available. Because of its light and penetrating nature very close attention must be paid to sealing the area to be treated.

Bin Surface Applications

Dilute with water to make enough spray to treat 1,000 ft.sq. of bin surface. Use only in empty bins.

- Storcide II (deltamethrin + chlorprifos-methyl) 1.8 fl oz in 1 gal.
- Tempo SC Ultra (cyfluthrin) 0.27 fl. oz.
- Insecto, etc₂ (silicon dioxide, from diatomaceous earth)0 lb. per 1,000 ft. sq.

Note: some grain buyers, especially food processors will not except grain coated with this product. Be sure your buyer will accept this treatment.

Grain Protectants

Applied directly to stored small grains.

Do not use the same compound for both Bin Surface and Grain protection.

	Amount per 1,000 bu.
Storcide II	12.4 fl. oz. -- Wheat
	9.9 fl. oz. -- Barley
	6.6 fl. oz. -- Oats

Grain Surface "Cap Out" Treatments

Applied directly to the top surface of stored small grains for Indian Meal Moth control.

Biobit HP (B.t.)	3 oz. / 1,000 ft. sq. (mixed to 4 inches deep)
Dipel DF (B.t.)	1 lb. / 1,000 ft. sq. (mixed to 4 inches deep)
Javelin WG (B.t.)	14 oz / 1,000 ft. sq. (mixed to 4 inches deep)

Indian meal moth larvae can be controlled by many products containing the active ingredient *Bacillus thuringiensis* "B.t.". Biobit, Dipel and Javelin are just examples of these products. B.t. products will not control beetles and weevils.

Note: Indian meal moth adults may be controlled by hanging DDVP Resin strips (Vapona) in the head space over the grain mass. Use 1 strip for each 1,000 cubic feet of air space over the grain. One treatment will last about 3 months.

Bulk Grain Fumigation

Amount of Fumigant to be applied/1,000 bu. stored small grains.

<i>Phostoxin, Fumtoxin etc.</i> (aluminum phosphide)	tablets 40 - 180 / 1,000 bu.
	pellets 200 - 900 / 1,000 bu.

Note: Economic thresholds are hard to determine for stored grain but these numbers should provide a guide to when fumigation will be profitable. Rice weevil or lesser grain borer 1 insect / qt of grain. Red flour beetle, rusty grain beetle and other bran bugs 5 insects / qt of grain. Successful fumigation includes consideration of many variables, use these fumigant amounts as guide and consult the label of the product you choose.

WARNING: Fumigation is a complicated and dangerous technique. If at all possible hire a commercial fumigator. If a commercial fumigation is not possible consult the label of the product you have chosen to use and follow it to the letter. Note that the Aluminum phosphide label has undergone major revision in recent years and now contains significant requirements for pre-planning and documentation of the fumigation and access to considerable safety equipment.

Information Summary Table for Small Grain Insecticides

This table is provided for a quick comparison of insecticides labeled on small grains. Insecticides are listed alphabetically by pesticide common name (usually present in the active ingredients section of the product label). One or more brand names are included along with the Restricted Entry Interval (REI) and Mode of Action Group number. Brand names of Restricted Use pesticides appear in *bold italics*.

Use pesticide products only in accordance with their labels and with the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval. Check the label for Personal Protective Equipment required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water.

Mode of Action Group A numerical classification system has been developed to make it easy to recognize the modes of action of insecticide products. Insecticides with the same mode of action belong to groups with unique numbers. Selection of a labeled product from a different number category (different mode of action) will help to slow down the development of resistance to either group. For example, alternate use of pyrethroid insecticides and pyrethrins sprays (Category 3) with labeled organophosphate insecticides (Category 1B). Always avoid tank mixing products with the same mode of action. These Mode of Action Group codes are on many pesticide labels and have been developed by the Insecticide Resistance Action Committee (IRAC).

Common Name	Brand Name	Restricted Entry Interval (hours)	Mode of Action Group
aluminum phosphide	<i>Phostoxin, Fumtoxin</i>	NA [‡]	8D
Bt kurstaki	Biobit, Dipel DF, Javelin WG, etc.	4	11B2
carbaryl	Sevin	12	1A
carbofuran	<i>Furadan 4F</i>	48	1A
chloropicrin	<i>Chlor-O-Pic</i>	NA [‡]	NA
chlorpyrifos-methyl	Storcide II	NA [‡]	1B
cyfluthrin	Tempo, Storcide, <i>Govern</i>	NA [‡]	3
<i>lambda</i> -Cyhalothrin	<i>Warrior, Mistic Z, Taiga Z</i>	24	3
<i>zeta</i> -Cypermethrin	<i>Mustang Max, Mustang</i>	12	3
deltamethrin	<i>Storcide II</i>	NA [‡]	3
dimethoate	Dimethoate	48	1B
disulfoton	<i>Di-Syston 15 G, 8E</i>	48*	1B
imidacloprid	Gaucho	12*	4A
malathion	Cythion, Malathion	12	1B
methyl bromide	<i>Bromo Gass</i>	NA [‡]	8 ^a
methomyl	<i>Lannate</i>	48	1A
methyl parathion	<i>Penncap-M</i>	4 days	1B
permethrin	<i>Ambush, Arctic</i>	12	3
Silicon Dioxide	Insecto	None listed	None listed
spinosad	Tracer	4	5

NA Not applicable in the usual sense.

* If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated.

‡ For use in storage bins no reentry is allowed. See label for details

Educational programs of the Kentucky Cooperative Extension Service serve all people regardless of race, color, age, sex, religion, disability, or national origin.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. M. Scott Smith, Director of Cooperative Extension Service, University of Kentucky College of Agriculture, Lexington, and Kentucky State University, Frankfort.

Revised annually — 10/09